

Phase Separation and Superconductivity in the Copper Oxide Chain.

C. Vermeulen¹, W. Barford¹ and E. R. Gagliano^{2*}

1.Department of Physics, The University of Sheffield, Sheffield, S3 7HR,
United Kingdom.

2.Physics Department, University of Illinois at Urbana-Champaign, 1110
W. Green, Urbana, IL 61801, U.S.A.

Abstract

The phase diagram of the copper-oxide chain as a function of density and the nearest-neighbour Coulomb interaction, V , is determined. Phase separation takes place above a critical value of V when the $\text{Cu}^{2+} \rightarrow \text{Cu}^+$ valence fluctuations dominate. In the proximity of the phase separation boundary the superconducting correlations are the most divergent. We identify the parameter regions where the Luttinger Liquid theory applies and calculate the contours of the charge-charge correlation exponent K_ρ . We show that anomalous flux quantization occurs as K_ρ diverges. At $n = 1$, for $V > t$, a gap opens in the spectrum and the ground state is a charge-density wave.

PACS numbers: 64.70, 74.10, 74.70V

*Present address: Centro Atomico Bariloche, 8400 Bariloche, Argentina.

The high temperature superconductors are difficult to understand quantitatively because of the failure of mean field theories and perturbation methods to give reliable results both in reduced dimensions and in the presence of strong electronic interactions. Under these circumstances exact calculations on finite size systems can provide valuable information. In particular, the results of Luttinger liquid theory enable thermodynamic correlation functions to be deduced for one dimensional systems via exact calculations on finite size chains. In this letter we solve exactly finite length copper-oxide chains, up to a maximum of 12 sites, using the Lanczos procedure with the anticipation that the copper-oxide plane will show some features in common with the copper-oxide chain [1]. Our purpose is to study the phase separation and superconducting instabilities arising from the nearest neighbour copper-oxygen Coulomb repulsion. We identify the phase separation boundary of the extended Hubbard model via a Maxwell construction. Then we identify the region of possible superconductivity using the results of Luttinger Liquid theory and by the onset of anomalous flux quantization.

Our model for the copper-oxide chain consists of two atoms per unit cell. Neglecting the oxygen-oxygen hybridization and considering the Coulomb interaction up to first-nearest neighbours, the copper-oxide chain is described by the two band model Hamiltonian,

$$H = H_t + \frac{\Delta}{2} \sum_{ij\sigma} (p_{j\sigma}^\dagger p_{j\sigma} - d_{i\sigma}^\dagger d_{i\sigma}) + U_d \sum_i d_{i\uparrow}^\dagger d_{i\uparrow} d_{i\downarrow}^\dagger d_{i\downarrow} + U_p \sum_j p_{j\uparrow}^\dagger p_{j\uparrow} p_{j\downarrow}^\dagger p_{j\downarrow} \\ + V \sum_{\langle ij \rangle \sigma \sigma'} (d_{i\sigma}^\dagger d_{i\sigma} p_{j\sigma'}^\dagger p_{j\sigma'}) + \left(\frac{(U_d + U_p)}{4} + 2V \right) (N_s - N_p),$$

where

$$H_t = -t \sum_{\langle ij \rangle \sigma} (d_{i\sigma}^\dagger p_{j\sigma} + h.c.). \quad (1)$$

i and j are copper and oxygen sites respectively, $\langle ij \rangle$ represents nearest neighbours and the operator $d_{i\sigma}^\dagger$ ($p_{j\sigma}^\dagger$) creates a Cu (O) hole with spin σ . Δ is the charge-transfer energy, U_d (U_p) is the copper (oxygen) Coulomb repulsion, N_s is the number of sites, N_p is the number of particles, and V and t are the copper-oxide Coulomb repulsion and hybridization, respectively. The Hamiltonian has been defined so that it is invariant under the ‘particle-hole transformation’ $d_{i\sigma}^\dagger \rightarrow d_{i\sigma}$, $p_{j\sigma}^\dagger \rightarrow p_{j\sigma}$ and $\Delta \rightarrow U_d - (\Delta + U_p)$, *i.e.* $E(N_p, \Delta) = E(2N_s - N_p, U_d - (\Delta + U_p))$.

In this letter we consider parameters which are chosen to fit, as far as possible, with the photoemission data of high temperature superconductors. Thus $U_d \sim 9t$, $\Delta \sim 2 - 3t$ and U_p small, where $t \sim 1.5eV$ [2]. Hence the region $0.5 < n < 1.0$ corresponds to $Cu^{2+} \rightarrow Cu^+$ valence fluctuations, while (because of the ‘particle-hole’ symmetry of equation (1)) the region $1.5 > n > 1.0$ corresponds to $Cu^{2+} \rightarrow Cu^{3+}$ valence fluctuations. As we are interested in the role played by the nearest neighbour interaction, V , this will be left as a free parameter.

For most regions of the phase diagram the system is metallic, for which Luttinger Liquid theory can provide a valuable insight. However, for some parameter regions in the doping range $0.5 < n < 1.0$ the system is unstable with respect to phase separation. This occurs at a finite value of the nearest neighbour Coulomb repulsion, and we identify this phase boundary before proceeding to a discussion of the metallic regime.

Figure 1 shows the phase separation boundary as a function of V and hole density for $U_p = 0$ and $U_p = t$. We set $U_d = 9t$ and $\Delta = 2t$; the number of sites used was 12. The boundary for this region is found using the Maxwell construction. This is given by the points (n_1, n_2) at which a tangent touches the curve of the ground state energy versus density. Other authors have used the point at which the compressibility, κ , diverges (*i.e.* the point at which the ground state energy versus density becomes concave). These points, however, actually lie between n_1 and n_2 and so could be a misleading indicator for when phase separation has actually occurred. Figure 1 also shows the line of $\kappa = \infty$. Using the Maxwell construction, the densities of the phase separated regions are then given by n_1 and n_2 , and the fraction of the system occupied by n_1 and n_2 at any given intermediate density n' is given by the relation $mn_1 + (1 - m)n_2 = n'$, where $m = \frac{(n_2 - n')}{(n_2 - n_1)}$. The tendency towards charge segregation can also be seen in the oxygen-oxygen charge correlation function [3], defined by

$$C_{cdw}(k) = \frac{1}{N_s} \sum_{ij} (n_i - \langle n \rangle)(n_j - \langle n \rangle) e^{ik(R_i - R_j)}, \quad (2)$$

where $n_i = \sum_{\sigma} c_{i\sigma}^{\dagger} c_{i\sigma}$.

The oxygen-oxygen charge correlation function (shown in figure 2) shows a peak at $k = \frac{\pi}{3}$ (*i.e.* one over the system length) for both a density of $n = \frac{8}{12}$ and $n = \frac{10}{12}$. This indicates a phase separation instability in the system. Phase separation occurs because, to avoid paying Coulomb repulsion, holes will tend to surround themselves with empty sites. This is because a repulsion, V , between a copper and a oxygen hole actually means an attraction, $-V$, between a copper hole and a oxygen electron. The result is that as the doping is increased away from $n = 0.5$ holes will tend to move onto those oxygen sites which are surrounded by empty copper sites. However, where possible holes will stay on copper sites to avoid paying the charge transfer

energy, Δ . The result is that the charge density becomes segregated and the Bloch symmetry breaks down [4].

The phase separated region is shifted as U_p is increased into higher values of V , as can be seen in the inset of figure 1. However, the phase separation boundaries do not depend strongly on U_p provided that $U_p < U_d - \Delta$ is satisfied.

Having determined the region of the phase diagram for which a phase separation instability occurs we now turn our attention to the metallic phase. In the metallic region the long distance behaviour of the correlation functions can be determined via the aid of Luttinger Liquid theory. Before discussing our results we summarise some of the important aspects of Luttinger Liquid theory.

Luttinger Liquids belong to the universality class of models described by conformal field theory with central charge, $c = 1$. This theory is useful because it gives relationships between the correlation exponents and a few simple properties of the finite system. Initially it was shown that models with only one degree of freedom, such as spinless fermions or bosons, had $c = 1$. Applying the same ideas to the Hubbard model proved difficult since conformal quantum field theory deals with Lorentz invariant systems only; *i.e.* it requires all gapless excitations to have the same velocity. For the Hubbard model this is true only at half filling, as then there is a gap for charge excitations.

Recently it has been shown that conformal field theory can be applied to models with more than one gapless excitation if the excitations are decoupled in the low energy regime. In the low energy regime this is true of the Hubbard model [5]. In such a case each degree of freedom has an associated operator algebra, which is again characterized by a central charge $c = 1$. Provided that there are no phase changes as the interactions are increased it is reasonable to assume that Luttinger Liquid theory applies to lattice quantum models in the strong coupling regime. In fact, this assumption can be tested *a posteriori* by showing that the predictions of Luttinger Liquid theory are internally consistent. This is the assumption adopted in this paper for the extended Hubbard model.

At long distances the charge-charge and spin-spin correlation functions are given by [6],

$$\langle n(x)n(0) \rangle = \frac{K_\rho}{(\pi x)^2} + A_1 \frac{\cos(2k_f x)}{x^{(1+K_\rho)}} \ln^{-3/2}(x) + A_2 \frac{\cos(4k_f x)}{x^{4K_\rho}}$$

and

$$\langle S_z(x)S_z(0) \rangle = \frac{1}{(\pi x)^2} + B_1 \frac{\cos(2k_f x)}{x^{(1+K_\rho)}} \ln^{1/2}(x),$$

respectively. The superconducting correlations are similar to the spin-spin correlator with K_ρ being replaced by $1/K_\rho$. Hence, K_ρ equals 1 for free

fermions with spin. $K_\rho < 1$ means that the charge and spin correlations dominate and $K_\rho > 1$ implies that the superconducting correlations are the most divergent.

Conformal field theory relates bulk quantities such as the compressibility, κ , and the Drude weight, D_c , with the charge-charge correlation exponent. Where the system behaves as a Luttinger Liquid, the exponent K_ρ is related to the compressibility by $K_\rho = \frac{\pi}{2}\kappa v_c n^2$ and to the Drude weight by $K_\rho = D_c \pi / v_c$ (where v_c is the charge velocity). When analytic solutions are not available these bulk quantities [7, 8] can be evaluated using exact diagonalization techniques [9]. The discrete compressibility is given by $N_p^2 \kappa = 4/[e(N_p + 2) + e(N_p - 2) - 2e(N_p)]$, where $e(N_p)$ is the ground state energy per site. To calculate the Drude weight the chain is threaded with a dimensionless constant flux, ϕ . This is done mathematically by the operator transformation, $c_{k\sigma}^\dagger \rightarrow c_{k\sigma}^\dagger e^{\frac{-ik\phi}{N_s}}$. The finite-size equivalent of the Drude weight is then $D_c = [e(\phi_0 + \delta\phi) - e(\phi_0)]/\delta\phi^2$, where $\delta\phi$ is an infinitesimal flux (~ 0.01) and ϕ_0 is a background flux which removes accidental degeneracies of the ground state and guarantees a *zero-current* ground state. For example, ϕ_0 is 0 (*periodic boundary conditions*) for $N_p = 4m + 2$ and $\frac{\pi}{N_s}$ (*anti-periodic boundary conditions*) for $N_p = 4m$ with m an integer [10].

The charge velocity, as well as the spin velocity, can also be obtained from finite cluster calculations of the response functions [11]. In fact, they are related to the lowest frequency pole ω_l of the charge-charge and spin-spin dynamical correlation functions by $v_l = \frac{\omega_l}{q_{min}}$, where $q_{min} = \frac{2\pi}{N_u}$, N_u is the number of unit cells and l refers to both charge and spin degrees of freedom [12]. One can therefore directly determine both velocities without adiabatically following the relevant low-energy excitations, as has recently been done for the $t - J$ model [7].

The phase diagram, figure 1, shows the contour of $K_\rho = 1$, and the region for which K_ρ exceeds 1. The expression used to calculate this contour was $K_\rho = \pi n \sqrt{\frac{D_c \kappa}{2}}$. Notice that for low doping K_ρ exceeds 1 within the phase separation region given by the Maxwell construction, and as doping is increased the curve $K_\rho = 1$ moves out indicating that the superconducting correlations begin to dominate well within the conducting region. The points on the contour corresponding to the densities $n = \frac{8}{12}$ and $n = \frac{10}{12}$ are found from the 12 site chain, the point at $n = \frac{8}{10}$ is found using a 10 site chain and the point at $n = \frac{6}{8}$ is found using an 8 site chain. At high densities the point of K_ρ exceeding 1 is associated with the appearance of anomalous flux quantization. Anomalous flux quantization is the condition that the ground state energy is periodic in ϕ , with period π , (*i.e.* the flux is quantised in units of $\frac{hc}{2e}$). For the attractive Hubbard model, which is known to exhibit singlet superconductivity, the ground state energy satisfies

$$E_0(\phi + \pi) - E_0(\phi) = \frac{\Lambda}{2} \left(\frac{N_s}{\zeta} \right)^{1/2} \exp \left(\frac{-N_s}{\zeta} \right), \quad (3)$$

where ζ is a length associated with a gap of spin excitations and Λ depends on details of the model [13]. As well as exhibiting anomalous flux quantization, the superconducting state is defined by the Drude weight (or superfluid density) remaining finite in the thermodynamic limit. For $n = \frac{10}{12}$ and $\frac{8}{10}$ anomalous flux quantization appears almost exactly at the point that K_ρ exceeds 1, the result being slightly better for $n = \frac{10}{12}$. For lower densities this relationship is not so clear.

We have also performed finite size scaling [14] using a 6 and 12 site chain at $n = \frac{2}{3}$. Using these two chains we are able to calculate the parameter ζ of equation (3) and the ratio $\frac{D_c(Ns=12)}{D_c(Ns=6)}$ for different values of V . We are then able to determine whether this is in the asymptotic region of equation (3) (*i.e.* the region of relatively large N_s and small ζ). A small value of the ratio of the Drude weights in the asymptotic region will indicate that $D_c = 0$ in the thermodynamic limit, and that the true ground state is non-conducting. Anomalous flux quantization, without conductivity, is the signal for a pairing but non-superconducting state, such as a phase separated state or a charge density wave. When $V = 4t$ (a point well within the phase separation region), $\zeta \sim 2$ so this is in the asymptotic region, but the ratio $\frac{D_c(Ns=12)}{D_c(Ns=6)} \sim 0.1$. Such a large reduction in Drude weight with such a small ζ indicates an insulating state. Anomalous flux quantization first occurs when $V \sim 2.7t$ for a 12 site chain at this doping. This indicates that at $V = 4t$ there is a definite phase separation instability.

For all other values of doping, namely $n < 0.5$ and $n > 1$, the effect of nearest neighbour repulsion is to suppress the superconducting correlations. For the densities $n < 0.5$ and $n > 1.5$, the phase diagram rather resembles the corresponding phase diagram of the $t - U - V$ model [8].

Finally, we consider the effect of the nearest neighbour repulsion at half filling, $n = 1$. At this filling the effect of increasing V is to force the system into an insulating phase. This can be seen from the graph of the Drude weight, the inset of figure 3, which vanishes at $V \sim 1.6t$. The driving force for this change is illustrated by the charge correlation function shown in figure 3. It shows that as V is increased there is a sharp rise in the correlation function at $k = \pi$ (a wavelength of *one* unit cell). This corresponds to charge being pushed from copper sites onto oxygen sites, resulting in an insulating charge density wave.

In conclusion, we have established the phase diagram of the copper-oxide chain using finite cluster calculations. In the region in which $Cu^{2+} \rightarrow Cu^+$ charge fluctuations dominate (*i.e.* $\Delta \ll U_d$) we find that the coupling to the nearest neighbour repulsion leads to a region of phase separation for a sufficiently large interaction (typically $V \sim \Delta$). In the proximity of the phase separation region and large enough density superconducting correlations are the most divergent. From our results we find that the curve $K_\rho = 1$ crosses the phase separation boundary at $n \sim 0.72$. At the densities $n = \frac{8}{10}$

and $\frac{10}{12}$ the divergence of K_ρ is accompanied by anomalous flux quantization. The contour $K_\rho = 1$ has been calculated near the metal-insulator transition ($n = 0.5$) with only a small number of doped holes. The trend for the superconducting region to disappear as $n \rightarrow 0.5$ may be a consequence of this fact. Close to the phase separation boundary the superconducting correlations are precursory to phase separation. At weaker coupling the role played by charge transfer excitons [15] is still to be understood. For all other parameter regions superconducting correlations are suppressed. At $n = 1$ the model shows a transition from a metallic state to a insulating charge density wave.

Acknowledgements

We thank the SERC (United Kingdom) for the provision of a Visiting Research Fellowship (ref. GR/H33091). W.B. also acknowledges support from the SERC (ref. GR/F75445) and a grant from the University of Sheffield research fund, while E.G. acknowledges support from DMR89-20538/24. C.J.V. is supported by a University of Sheffield scholarship. We thank B. Alascio, A. Aligia, S. Bacci, C. Balseiro, E. Fradkin, G. A. Gehring and R. M. Martin for useful conversations. The computer calculations were performed at the National Center for Supercomputing Applications, Urbana, Illinois and at the University of Sheffield.

Bibliography

- [1] These studies are also interesting in their own right, however, because copper-oxide chains do exist in some of the high temperature superconductors, for example $\text{YBa}_2\text{Cu}_3\text{O}_7$ (in the form of copper-oxide diamonds). Furthermore, the one dimensional halogen-bridged metal complexes are also charge transfer systems which exhibit many of the fascinating features of polyacetylene, for example charge density waves, spin density waves and solitons. To model true one dimensional systems, however, electron-phonon couplings should be included to account for the Peierls transition. This will not be done in this paper.
- [2] G. A. Sawatzky, 1991. High Temperature Superconductivity. ed. D. P. Tunstall and W. Barford, Publishers: Adam Hilger (Bristol).
- [3] K. Sano and Y. Ono, Phys. C. **205**, 170 (1993).
- [4] W. Barford and M. W. Long, J. Phys. Condens. Matt., **5**, 199 (1993)
- [5] H. Frahm and V. E. Korepin, Phys. Rev. B, **42**, 10553 (1990).
- [6] H. J. Schulz, Phys. Rev. Lett., **64**, 2831 (1990).
- [7] M. Ogata, M. U. Luchini, S. Sorella and F. F. Assaad, Phys. Rev. Lett., **66**, 2388 (1991).
- [8] H-Q.Lin *et al.*, submitted 1993 and references therein.
- [9] E. Gagliano, E. Dagotto, A. Moreo and F. C. Alcaraz, Phys. Rev. B, **34**, 1677 (1986).
- [10] For a two-band spinless Fermion model, antiperiodic boundary conditions are required at any density in order to have a zero current ground state.
- [11] E. Gagliano and C. Balseiro, Phys. Rev. Lett., **59**, 2999 (1987) and, Rev. B. **43**, 13660 (1991).
- [12] Since the charge operator does not conserve total spin, the peak associated with v_s , appears also in the charge-charge response function. Hence, the charge velocity is given by the low energy peak that does not appear simultaneously in both dynamical correlations.

- [13] C. A. Stafford *et al.*, Phys. Rev. B **43**, 13660 (1991)
- [14] A. Sudbo et al, Phys. Rev. Lett. **70**, 978 (1993).
- [15] C. J. Vermeulen, W. Barford and E. R. Gagliano, submitted to Phys. Rev. Lett., 1994

Figure 1: The phase diagram of the copper-oxide chain as a function of V and hole density for $U_d = 9t$, $U_p = 0$ and $\Delta = 2t$, showing the phase separation boundary, the contour of $K_\rho=1$ and the contour of infinite compressibility. There is also shown the onset of anomalous flux quantization for $n = \frac{10}{12}$ and $\frac{8}{10}$. The inset shows the phase separation boundary for $U_p = 0$ and t .

Figure 2: The oxygen-oxygen charge correlation function for $n = \frac{10}{12}$ and $\frac{8}{12}$. This shows a peak at $k = \pi/3$ for $V = 2t$, indicating a phase separated state ($U_d = 9t$, $U_p = 0$ and $\Delta = 2t$)

Figure 3: The Drude weight as a function of hole density for $U_d = 9t$, $U_p = 0$ and $\Delta = 2t$, showing the opening of the gap at $n = 1$. The copper-oxygen charge-charge correlation function for a 12 site chain at $n = 1$ is shown for $V = 0.5t$ and $2t$. The density of electrons on the oxygen atoms rises from $n = 1.265$ to $n = 1.797$ in this range.